

Advanced Cardiac Life Support

Advanced Provider Overview

Contact Information

- Judy Haluka, BS, RCIS, EMT-P
- Emsconed.org
- Jhaluka@emsconed.org
- Fax: 412-299-2026
- Regional Faculty – American Heart Association
 - ACLS – PALS - BCLS

References

- Currents in Emergency Cardiovascular Care(Winter 2005-2006)
- Circulation 2005;112:IV-1-IV-5) © American Heart Association

Basic Life Support

- Importance of Coronary Perfusion Pressure
- 30:2 Compression/Ventilation Ratio
- Complete Chest Recoil
- No Interruptions
- No Hyperventilation (6-8 without perfusion: 10-12 with perfusion)

Basic Cardiac Life Support

- Advanced Airway
 - 100 Compressions/min
 - 6-8 Ventilations/min
 - Push fast – push hard
- Switch Providers at 2 Minutes

Successful Resuscitation

- Survival to discharge from facility with near normal neurological function
 - 10 minute recommendation
 - 30 minute guideline
 - Excluding special circumstances
- Worked to pronouncement or ROSC – no transport with CPR

Primary Survey

- Airway
- Breathing
- Circulation
- Defibrillation

Secondary Survey

- Airway – Adequate/Obstructed
 - Determine if advanced airway needed
- Breathing – confirm airway placement
 - Pneumothorax, flail chest, open chest wound
 - Air movement - oxygenation

Secondary Survey

- Circulation
 - IV access
 - Monitor – determine rhythm and required treatment
 - Fluids – if indicated

Secondary Survey

- Differential Diagnosis
 - Determine cause of arrest or pre-arrest state
 - Institute definitive care based upon diagnosis

Secondary Survey - Advanced

- Disability
 - Mental status (glasgow coma scale – stroke scale if indicated)
- Expose
 - Completely expose patient, check for injuries, lesions, temperature
 - Check extremity pulses

Secondary Survey - Advanced

- Fingers – Foley – Flip
 - Rectal, vaginal exam
 - Pelvic injuries
 - Flip patient to check back
 - Identify evidence of trauma, pregnancy, sources of bleeding

Secondary Survey - Advanced

- Gastric Tube
 - Check aspirate for blood, ingested tablets, toxins
- History
 - Document
 - Question family, friends, EMS personnel

Quadrant Three

- Oxygen – Monitor – IV – Fluids
 - Identify hypoxia, symptomatic arrhythmias, hypovolemia, treat and evaluate response to therapy

Quadrad Four

- Temperature – Heart Rate – Blood Pressure – Respirations
 - Support as needed, consider vasoactive therapy; evaluate response and adjust therapy

Quadrant Five

- Tank (volume)
- Tank (resistance)
- Pump (Failure)
- Rate

Ventricular Fibrillation

- Witnessed – immediate defibrillation
- Unwitnessed – CPR 2 minutes
- Defibrillation
 - 360 joules if monophasic
 - 200 joules – biphasic
 - One Defibrillation Only – NO MORE STACKED SHOCKS

Ventricular Fibrillation

- 2 minutes of CPR – no rhythm or pulse check
- Epinephrine 1mg q 3-5 minutes
- Vasopressin 40 units (single dose)
- Defibrillation

Ventricular Fibrillation

- Amiodarone 300mg IV (can be repeated once at ½ dose)
- If unavailable – Lidocaine 1 to 1.5mg/kg; then half dose up to a maximum of 3 doses
- Defibrillation

Ventricular Fibrillation

- Consider Magnesium 1 to 2 grams
 - If prolonged QT interval with Torsades de Pointe arrest
 - Not routine administration

Ventricular Fibrillation

- Precordial Thump – not indicated
- Sodium Bicarbonate – no longer considered until ROSC with gases or special circumstances
- No routine fluid bolus

PEA/Asystole

- Asystole – confirmation of death; absent special circumstances – not survivable
- Pulseless Electrical Activity
 - True EMD
 - Psuedo EMD

PEA/Asystole

- 2 Minutes of CPR
- Vasopressor
 - Epinephrine 1mg q 3-5 minutes
 - Vasopressin 40 units IV Push
- Atropine 1mg IVP if Asystole or Bradycardia
- Fluid Bolus if Tachycardia

PEA Differential

- Hypovolemia
- Hypoxia
- Hydrogen Ion (Acidosis)
- Hypo/Hyperkalemia
- Hypothermia

PEA Differential

- Toxins
- Tamponade
- Tension Pneumothorax
- Thrombosis (Coronary or Pulmonary)
- Trauma

Bradycardia

- Arrhythmias Don't Kill – Heart Rate Does
- Assess for Adequate Perfusion
 - If adequate – monitor
 - If inadequate – intervene
 - Goal is increase in heart rate
 - Conversion is secondary

Bradycardia

- Atropine 0.5mg (to total of 3mg)
 - Note change in dosage
- Transcutaneous pacing
 - First line for High Degree AV blocks (Mobitz II and Third Degree)
- Consider Epinephrine (2-10mcgms/min) or Dopamine (2-10mcgms/min)
 - As a Bridge to definitive intervention only

Tachycardia - Unstable

- Narrow or Wide Complex
 - Immediate Cardioversion

Tachycardia – Narrow Complex

- Stable
- Regular Rhythm
 - Attempt Vagal Maneuvers
 - Adenosine 6mg IVP, 12mg, 12mg
 - Stopcock method
 - Extremely short half life

Tachycardia – Narrow Complex – Regular Rhythm

- Conversion with Adenosine
 - Probable Reentry SVT
 - Observe for recurrence
 - Treat recurrence with Adenosine
 - Long acting AV nodal blocking agent
 - Diltiazem, Beta Blockers

Tachycardia – Narrow Complex

- Does not Convert with Adenosine
- Possible Atrial Flutter, Atrial Tachycardia or Junctional Tachycardia
 - Control Rate (Diltiazem, Beta Blockers)
 - Treat underlying cause
 - Consider EP Consult

Tachycardia – Narrow Complex Irregular Rhythm

- Probable Atrial Fibrillation, Flutter or MAT
 - Consider EP consult
 - Control Rate with Diltiazem, Beta Blockers

Tachycardia – Wide Complex Stable

- Regular Rhythm
 - Probable Ventricular Tachycardia
 - Treat with Amioderone 150mg IV over 10 minutes (repeat as needed to maximum of 2.2 grams in 24 hours)
 - Prepare for elective cardioversion
 - If SVT with Aberrance – treat with Adenosine

Tachycardia – Wide Complex Irregular Rhythm

- Probable Atrial Fibrillation with Aberrancy
 - Treat as if narrow complex tachycardia
- If pre-excited Atrial Fibrillation (AF + WPW)
 - EP Consult Advised
 - Avoid AV Nodal Blockers (Adenosine, Verpamil, Diltiazem, Digoxin)

Tachycardia – Wide Complex Irregular Rhythm

- Consider Amiodarone 150mg IV over 10 minutes
- If Torsades de Pointe, consider Magnesium 1-2 grams over 5 – 60 minutes.

Acute Coronary Syndromes

- ST Segment Elevation or new LBBB with Symptoms of AMI
 - High specificity for evolving STEMI; assess for reperfusion candidacy
- ST Depression
 - Consistent with strong suggestion of ischemia; high risk subset of patients with unstable angina

Acute Coronary Syndromes

- Non-diagnostic ECG
 - Further assessment required
 - Serial enzymes; repeat ECG
 - Consider other causes of chest pain: aortic dissection, pericarditis/myocarditis, pulmonary embolism

AMI Reperfusion Guidelines

- ECG within 10 minutes
- Door to drug time less than 30 minutes
- Door to balloon time less than 90 minutes

AMI Protocol

- Oxygen – IV Access – Continuous ECG Monitoring
- Reperfusion Therapy (PCI or fibrinolytics) for STEMI
- Prompt Aspirin – 160mg to 325mg for all patients
- Beta blockade for all patients without contraindication
- IV NTG for initial 24-48 hours with AMI and CHF, large AWTMI, persistent ischemia or hypertension

Potential Adjunctive Therapy

- Beta Blockers
 - Decreases myocardial oxygen consumption
 - Increases myocardial salvage in area of infarction
 - Can reduce VT and fibrillation
 - Benefits: 23% reduction in long term mortality

Heparin

- All patients undergoing PCI or surgical revascularization
- All patient receiving fibrinolytics
- IV or SQ low molecular weight in patients with NSTEMI
- IV unfractionated heparin for patients at increased risk of systemic emboli (atrial fib, large anterior MI, LV thrombus)

ACE Inhibitors

- Early oral therapy reduces mortality and CHF associated with AMI
- Helps prevent adverse LV remodeling
- Delays progression of heart failure
- Decreases sudden death
- Decreases recurrent MI

Cardiac Markers

- Myoglobin 1-2 hours
- Cardiac troponins 3-12 hours
- CK-MB 3-12 hours

NSTEMI – Adjunctive Therapy

- Low molecular Weight Heparin
 - Inhibits thrombin
 - Predictable results
- IIb/IIIa Inhibitors
 - Reduction in death and MI
 - Major benefits realized when PCI is planned

NSTEMI – Adjunctive Therapy

- Plavix
 - 300mg loading dose in addition to aspirin
 - Can substitute for ASA if true allergy
- Statins
 - Reduces incidence of reinfarction, recurrent angina, rehospitalization and stroke when administered within a few days of symptom onset

Airway Management

- Advanced airway (intubation) may be deferred until ROSC
 - Particularly if Anesthesia Personnel not available
 - Increased complication when intubated by non-anesthesia personnel
 - Complications directly related to number of intubations performed